

| ChemoNavi | | | 投与 サイクル 数 | 1コース | 治療内容 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|------------|-----------------|---|---|--|---|---------------|----|-----|----|----|----|----|----|----|----|----|----|----|----|-----|----|----|----|-----|----|----|----------------|---|---|---|---|--|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|----|----|----|--|----------------|---|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|----------------|---|---|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|--|-----------------|---|--|--|--|--|---|--|--|--|--|--|--|---|--|--|--|--|--|--|--|---|--|--|--|-----------------|--|--|--|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|--------------|----------|--|
| 薬品名※()は成分名 | 投与量 | 投与日 | | | day 1 | day 2,3 | day 8, 15, 22 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| プレドニン(プレドニゾン)*1 | 45mg/㎡ | day1~21 | 総投与時間 day1:4時間30分 day2,3:1時間20分 day8,15,22:30分 | 点滴静注 ① 生理食塩液 100mL 10:00~ ルート確保・フラッシュ用 ↓ ② グラニセトロン 静注バッグ 50mL 10:10~ 全開 ↓ ③ オンコピン 50mL mg 生理食塩液 50mL 10:20~ 全開 ↓ ④ ダウノマイシン 50mL mg 生理食塩液 100mL 10:30~11:30 1時間 ↓ ⑤ エンドキサン 500mL mg 生理食塩液 500mL 11:30~14:30 3時間 | 点滴静注 ① 生理食塩液 100mL 10:00~ ルート確保・フラッシュ用 ↓ ② グラニセトロン 静注バッグ 50mL 10:10~ 全開 ↓ ③ ダウノマイシン 50mL mg 生理食塩液 100mL 10:20~11:20 1時間 | 点滴静注 ① 生理食塩液 100mL 10:00~ ルート確保・フラッシュ用 ↓ ② グラニセトロン 静注バッグ 50mL 10:10~ 全開 ↓ ③ オンコピン 50mL mg 生理食塩液 50mL 10:20~ 全開 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| エンドキサン(シクロホスファミド) | 900mg/㎡ | day1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ダウノマイシン(ダウノルピジン) | 30mg/㎡ | day1~3 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| オンコピン(ピンクリスチン)*2 | 1.3mg/㎡ | day1, 8, 15, 22 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| スプリセル(ダサチニブ) | 100mg/body | day4~31 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| *1 Day22~28日に漸減、終了。 *2 最大量2mg/body 寛解導入1は内服でステロイド、ダサチニブを使用。 | | | 血管外漏 出リスク | エンドキサン:炎症性 ダウノマイシン:壊死性 オンコピン:壊死性 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 投与スケジュール | | | 投与時の 注意事項 | day 1~21 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| JALSG-Ph+ALL213寛解導入療法2(60歳以上) <table border="1"> <thead> <tr> <th>治療日</th> <th>1</th> <th>2</th> <th>3</th> <th>4</th> <th>...</th> <th>8</th> <th>9</th> <th>10</th> <th>11</th> <th>12</th> <th>13</th> <th>14</th> <th>15</th> <th>16</th> <th>17</th> <th>18</th> <th>19</th> <th>20</th> <th>21</th> <th>22</th> <th>...</th> <th>28</th> <th>31</th> </tr> </thead> <tbody> <tr> <td>プレドニン 45mg/㎡*1</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td></td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>*1</td> <td>*1</td> <td>*1</td> <td></td> </tr> <tr> <td>エンドキサン 900mg/㎡</td> <td>●</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>ダウノマイシン 30mg/㎡</td> <td>●</td> <td>●</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>オンコピン 1.3mg/㎡*2</td> <td>●</td> <td></td> <td></td> <td></td> <td></td> <td>●</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>●</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>●</td> <td></td> <td></td> <td></td> </tr> <tr> <td>スプリセル100mg/body</td> <td></td> <td></td> <td></td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> <td>●</td> </tr> </tbody> </table> *1 day22~28日に漸減、終了 *2 最大量2mg/body | | | | 治療日 | 1 | 2 | 3 | 4 | ... | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | ... | 28 | 31 | プレドニン 45mg/㎡*1 | ● | ● | ● | ● | | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | *1 | *1 | *1 | | エンドキサン 900mg/㎡ | ● | | | | | | | | | | | | | | | | | | | | | | | | ダウノマイシン 30mg/㎡ | ● | ● | | | | | | | | | | | | | | | | | | | | | | | オンコピン 1.3mg/㎡*2 | ● | | | | | ● | | | | | | | ● | | | | | | | | ● | | | | スプリセル100mg/body | | | | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | 投与時の 注意事項 | day 4~31 | |
| 治療日 | 1 | 2 | 3 | 4 | ... | 8 | 9 | 10 | 11 | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 | ... | 28 | 31 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| プレドニン 45mg/㎡*1 | ● | ● | ● | ● | | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | *1 | *1 | *1 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| エンドキサン 900mg/㎡ | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| ダウノマイシン 30mg/㎡ | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| オンコピン 1.3mg/㎡*2 | ● | | | | | ● | | | | | | | ● | | | | | | | | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| スプリセル100mg/body | | | | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | ● | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| *1 day22~28日に漸減、終了 *2 最大量2mg/body | | | 備考欄 | 寛解導入療法2のDay1に 中枢神経予防として メソ レキセート15mg+デキサ メタゾン3.3mgの髄腔内注 射を考慮 | 内服 プレドニン 錠 分2 朝昼 | 内服 スプリセル 錠 分1 朝 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |